

## DISCUSSING IBS-D\*

\*IBS-D: irritable bowel syndrome with diarrhea

## Navigating Conversations About Your IBS-D Symptoms

Discussing your symptoms with your healthcare provider (HCP\*) might be uncomfortable, but it's an essential step to finding a treatment plan that is best for you. Use this conversation guide to help communicate your experience with your HCP and cover any questions you may have.

\*Healthcare provider may be a doctor, physician's assistant, or nurse practitioner

| Information For Your HCP  Before your appointment, please review this guide for your discussion with your HCP and print it out to take notes on the front and back.  |  |
|--|--|
| l've experienced the following symptoms in the last 12 mo (please check all that apply):  ☐ Abdominal Pain ☐ Diarrhea ☐ Bloating ☐ Othe  | I have/have not been diagnosed with  |
| Please fill out the information below to discurrence fill out the in | and diarrhea could be IBS-D  y anemia of colorectal disease, IBD bowel disease)  and diarrhea could be IBS-D I would/would not like to discuss if Xifaxan may be right for me  |
| Questions to Ask Your HCP About XIFAXAN  |  |
| Could XIFAXAN be right for me?     My questions:   |  |
| <ul> <li>INDICATION</li> <li>XIFAXAN® (rifaximin) 550 mg tablets are indicated for the treatment of irritable bowel syndrome with diarrhe (IBS-D) in adults.</li> <li>IMPORTANT SAFETY INFORMATION</li> <li>XIFAXAN is not for everyone. Do not take XIFAXAN if you have a known hypersensitivity to rifaximin, any of the rifamycin antimicrobial agents, or any of the components in XIFAXAN.</li> <li>If you take antibiotics, like XIFAXAN, there is a change</li> </ul>   | <ul> <li>XIFAXAN may affect warfarin activity when taken together. Tell your healthcare provider if you are taking warfarin because the dose of warfarin may need to be adjusted to maintain proper blood-thinning effect.</li> <li>If you are pregnant, planning to become pregnant, or nursing, talk to your healthcare provider before taking XIFAXAN because XIFAXAN may cause harm to an</li> </ul> |

does not improve or worsens. Talk to your healthcare provider before taking XIFAXAN if you have severe hepatic (liver) impairment, as this may cause increased effects of the medicine.

you could experience diarrhea caused by an overgrowth

ranging in severity from mild diarrhea to life-threatening

colitis. Contact your healthcare provider if your diarrhea

of bacteria (C. difficile). This can cause symptoms

• Tell your healthcare provider if you are taking drugs called P-glycoprotein and/or OATPs inhibitors (such as cyclosporine) because using these drugs with XIFAXAN may lead to an increase in the amount of XIFAXAN absorbed by your body.

unborn baby or nursing infant.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/ medwatch/ or call 1-800-FDA-1088.

For product information, adverse event reports, and product complaint reports, please contact: Salix Product Information Call Center Phone: 1-800-321-4576 | Fax: 1-510-595-8183 Email: salixmc@dlss.com

Please see accompanying <u>full Prescribing Information</u>.

